

## Systematic Review

# Thyroid Collision Tumors: A Systematic Review

Ari M. Abdullah<sup>1,2</sup>, Aras J. Qaradakhly<sup>1,3</sup>, Rawa M. Ali<sup>1,4</sup>, Rebaz M. Ali<sup>1,5</sup>, Yousif M. Mahmood<sup>1</sup>, Sami S. Omar<sup>6,7</sup>, Hawkar A. Nasralla<sup>1</sup>, Aso S. Muhiyaldeen<sup>1</sup>, Yadgar A. Saeed<sup>1</sup>, Hardi M. Dhair<sup>4</sup>, Rebaz O. Mohammed<sup>1</sup>, Hiwa O. Baba<sup>1,7</sup>, Berun A. Abdalla<sup>8</sup>, Fahmi H. Kakamad<sup>9</sup>, Abdulwahid M. Salih<sup>9\*</sup>

1. Smart Health Tower, Madam Mitterrand Street, Sulaymaniyah, Kurdistan, Iraq
2. Department of Pathology, Sulaymaniyah Teaching Hospital, Sulaymaniyah, Kurdistan, Iraq
3. Department of Radiology, Shorsh Teaching Hospital, Sulaymaniyah, Kurdistan, Iraq
4. Hospital for Treatment of Victims of Chemical Weapons, Halabja, Kurdistan, Iraq
5. Hiwa Cancer Hospital, Sulaimani Directorate of Health, Sulaymaniyah, Kurdistan, Iraq
6. Rizgary Oncology Center, Peshawa Qazi Street, Erbil, Kurdistan, Iraq
7. Kscien Organization for Scientific Research (Middle East Office), Hamid Street, Azadi Mall, Sulaymaniyah, Kurdistan, Iraq
8. Department of Biology, College of Education, University of Sulaimani, Madam Mitterrand Street, Sulaymaniyah, Kurdistan, Iraq
9. College of Medicine, University of Sulaimani, Madam Mitterrand Street, Sulaymaniyah, Kurdistan, Iraq

\* Corresponding author: [abdulwahid.salih@univsul.edu.iq](mailto:abdulwahid.salih@univsul.edu.iq) (A.M. Salih). Ablagh, House number 80, Zip code: 46001, Sulaymaniyah, Iraq



Keywords:  
PTC and MTC  
PTC and FTA  
Recurrence  
Collision

Received: April 20, 2024  
Revised: May 30, 2024  
Accepted: June 22, 2024  
First Published: June 28, 2024

Copyright: © 2024 Abdullah et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Abdullah AM, Qaradakhly AJ, Ali RM, Ali RM, Mahmood YM, Omar SS, et al. Thyroid Collision Tumors: A Systematic Review. Barw Medical Journal. 2024;2(3):44-56. <https://doi.org/10.58742/bmj.v2i2.94>

## Abstract

### Introduction

Collision tumors in the thyroid gland are exceedingly uncommon, comprising approximately 1.0% of all thyroid malignancies. This study aims to systematically review the presentation and management of collision tumors of the thyroid gland.

### Methods

A systematic review of published studies on thyroid collision tumors was conducted. All studies concerning thyroid collision tumors meeting the following criteria were included: 1) Confirmation of collision tumors via diagnostic methods, surgical exploration, or histopathological examination. 2) Presentation of case details within the study. 3) Tumors located precisely in the thyroid gland.

### Results

In total, 57 studies were compatible with the inclusion criteria. Most cases were female 87 (71.31%), and the remaining 35 (28.69%) were male. The patients' ages were between 12 and 88 years old, with a mean of  $49.87 \pm 14.48$  years. The most commonly presented symptom was neck swelling (45.08%). The most prevalent surgical procedure observed was total thyroidectomy combined with lymph node dissection, performed in 40 cases (32.78%). The predominant histopathological findings consisted of the simultaneous presence of papillary thyroid carcinoma and medullary thyroid carcinoma, identified in 51 cases (41.80%). The recurrence rate was observed in only 10 cases (8.20%).

### Conclusion

Thyroid collision tumors may primarily affect females, with the most frequent collisions being papillary and medullary thyroid carcinoma. Total thyroidectomy with lymph node dissection is the prevalent management option, and the recurrence rate can be lower than 10%.

## 1. Introduction

Collision tumors represent a phenomenon wherein two histologically distinct tumors, characterized by differing cellular lineages and genetic origins, coexist within the same mass and organ without a discernible transitional zone. This phenomenon frequently manifests in the liver, stomach, adrenal gland, ovary, lungs, kidneys, and colon [1,2].

Collision tumors can arise within the same organ, like renal cell carcinoma alongside transitional cell carcinoma, hepatocellular carcinoma with cholangiocarcinoma, and gastric adenocarcinoma with lymphoma. They can also occur in neighboring organs, such as sigmoid adenocarcinoma with urinary bladder transitional carcinoma. Additionally, they may coincide with systemic malignancies, like renal cell carcinoma with intravascular lymphomatosis, or occur as metastatic occurrences, such as breast carcinoma spreading to meningioma [3]. However, thyroid carcinoma stands as the predominant form of endocrine malignancy, accounting for 2.1% of cases, collision tumors in the thyroid gland are exceedingly uncommon, comprising approximately 1.0% of all thyroid malignancies, most of which are made up of papillary and medullary carcinoma [1,4-6].

Nodules form when growth signals intensify, prompting hyperplasia, or upon acquisition of a new genetic mutation, resulting in self-directed growth. Clinicians focus on discerning a thyroid nodule's benign or malignant nature during evaluation. The preferred diagnostic method is fine needle aspiration biopsy (FNAB). Notably, most thyroid nodules are benign, with malignancy detected in only 9-13% of cases [7,8].

In differentiated thyroid cancers, FNAB is often effective in diagnosing papillary carcinoma. However, its utility is restricted when diagnosing follicular and Hurthle cell tumors. This limitation stems from the fact that distinguishing between adenomas and carcinomas in these cases necessitates evidence of capsule invasion, which can only be detected through histological examination [9]. A frequently observed occurrence among thyroid collision tumors involves the pairing of papillary thyroid carcinoma (PTC) with medullary thyroid carcinoma (MTC) [2].

Managing thyroid collision tumors presents difficulties because of the differences in biological aggressiveness, treatment options, and prognosis among the tumors involved [10]. Decisions regarding treatment plans, such as surgical removal, radioactive iodine (RAI) therapy, chemotherapy, external radiotherapy, and additional supportive treatments, should be tailored to each patient based on the aggressive nature of the tumors involved, as this will ultimately influence the patient's prognosis [11].

This study aims to systematically review the presentation and management of collision tumors of the thyroid gland.

## 2. Methods

### 2.1. Study design

The current systematic review conformed to the guidelines outlined in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA).

### 2.2. Data sources and search strategy

A systematic review of all published studies on thyroid collision tumors was conducted utilizing Google Scholar and PubMed. The search incorporated specific keywords: (synchronous OR simultaneous OR collision thyroid cancer OR thyroid tumor OR thyroid carcinoma) AND (thyroid coexisting cancer OR tumor).

### 2.3. Eligibility criteria

Non-English-language studies and those unrelated to humans were excluded before or during the initial screening process. All studies concerning thyroid collision tumors meeting the following criteria were included: 1) Confirmation of collision tumors via diagnostic methods, surgical exploration, or histopathological examination. 2) Presentation of case details within the study. 3) Tumors located precisely in the thyroid gland. Studies published in predatory journals (lacking appropriate peer review) [12], as well as those not meeting the inclusion criteria, were excluded.

### 2.4. Study selection process

Initially, the titles and abstracts of identified studies underwent a preliminary screening, followed by a thorough examination of the full texts to determine eligibility. Various data of information were documented from the selected studies, including study design, participants' age and gender, symptoms, laboratory findings, diagnosis, type and site of the malignancies, treatment approaches, administration of RAI, outcomes, and recurrence rates.

### 2.5. Data item

The gathered data underwent analysis using Statistical Package for the Social Sciences software version 25.0 and were subsequently displayed in terms of frequency and percentages.

## 3. Results

In total, 340 studies were obtained from the resources, 19 of which were removed before any screening due to duplication, non-English language, non-article, and only abstracts. On the initial screening, the titles and abstracts of 247 studies did not match the inclusion criteria, and they were excluded. Overall, 74 studies underwent full-text screening, and 61 were assessed for eligibility. Finally, 57 studies [1- 3, 5- 7, 10, 11, 13- 61] (122 cases) were compatible with the inclusion criteria (Figure 1) (Table 1). Of the included studies, 50 (87.72%) were case reports, and 7 (12.28%) were case series. Most cases were female 87 (71.31%), and the remaining 35 (28.69%) were male. The patients' ages were between 12 and 88 years old, with a mean of  $49.87 \pm 14.48$  years. The most commonly presented symptom was neck swelling (45.08%). Available preoperative




Figure 1. Study selection PRISMA flow chart.

laboratory examinations revealed that thyroid-stimulating hormone (TSH) levels were within the normal range in 40 cases (32.79%), while thyroglobulin (TG) levels were elevated in 12 cases (9.84%), and calcitonin levels were elevated in 20 cases (16.39%). The most prevalent surgical procedure observed was total thyroidectomy combined with lymph node dissection, performed in 40 cases (32.78%). Moreover, cervical lymph node metastasis was present in 46 cases (37.70%). The predominant histopathological findings consisted of the simultaneous presence of PTC and MTC, identified in 51 cases (41.80%). This was followed by instances of PTC coexisting with follicular thyroid carcinoma, observed in 24 cases (19.67%). The isthmus exhibited the lowest frequency of malignancies, accounting for 7 cases (5.73%). Radioactive iodine was administered to 44 patients (36.06%). The recurrence rate was observed in only 10 cases (8.20%) (Table 2) (Table 3).

#### 4. Discussion

Collision tumors denote the coexistence of two histologically distinct neoplasms within the same anatomical location without intermixing. While the incidence of collision tumors is notably rare across all body regions, their occurrence in the thyroid gland is even more infrequent [3].

The terminology collision tumors of the thyroid have been utilized interchangeably with mixed and composite tumors. These designations denote the presence of multiple synchronous tumors exhibiting parafollicular and follicular-derived cellular elements within the thyroid gland. However, it is crucial to note that these terms are not synonymous, as they denote distinct entities. Within the context of thyroid tumors, a "mixed tumor" is exemplified by the Mixed Medullary and Follicular Tumor of Thyroid. This entity is a malignant tumor demonstrating morphological and immunophenotypical evidence of the coexistence of follicular and parafollicular tumor cells closely admixed within the same lesion [53]. Composite tumors often emerge from a common driver mutation in a neoplastic source, resulting in divergent histology [57]. Conversely, collision tumors represent histologically distinct tumors situated nearby within the same organ [53,62].

Several hypotheses have been advanced to elucidate the occurrence of collision tumors. The chance theory postulates the







**Table 2.** Basic characteristics summary of the included studies.

Variables	Frequency / Percentage
Age (mean of means) ± SD	49.87 ± 14.48
Gender	
Female	87 (71.31%)
Male	35 (28.69%)
Pre-operation Investigations	
TSH	
Low	6 (4.92%)
Elevated	3 (2.46%)
Normal	40 (32.79%)
N/A	73 (59.83%)
Thyroglobulin	
Low	0 (0%)
Elevated	12 (9.84%)
Normal	5 (4.10%)
N/A	105 (86.06%)
Calcitonin	
Low	0 (0%)
Elevated	20 (16.39%)
Normal	3 (2.46%)
Undetectable	3 (2.46%)
N/A	96 (78.68%)
A-TPO	
Low	0 (0%)
Elevated	2 (1.64%)
Normal	1 (0.82%)
N/A	119 (97.54%)
Presentation	
Neck Swelling	55 (45.08%)
Neck Pain	7 (5.73%)
N/A	64 (44.26%)
Histopathological Findings	
PTC and MTC	51 (41.80%)
PTC and FTC	24 (19.67%)
PTC and FTA	8 (6.55%)
PTMC and FTC	7 (5.74%)
PTC and HCA	5 (4.10%)
MTC, FTA, and PTMC	3 (2.46%)
PTMC and HCA	3 (2.46%)
PTC and SCC	3 (2.46%)
PTC and HCC	2 (1.64%)
PTC and MALT lymphoma	2 (1.64%)
MTC and FTA	1 (0.82%)
MTC and Melanoma	1 (0.82%)
OC, HPTC, and PDTC	1 (0.82%)
OC and PTMC	1 (0.82%)
Osteosarcoma and PTC	1 (0.82%)
PTC and adenocarcinoma	1 (0.82%)

PTC and MEC	1 (0.82%)
PTC and Liposarcoma	1 (0.82%)
PTC and TA	1 (0.82%)
PTC, FTC, and anaplastic	1 (0.82%)
PTMC and Anaplastic	1 (0.82%)
TVC and HCC	1 (0.82%)
FTC and NIFTP	1 (0.82%)
PTC, FTC, and MTC	1 (0.82%)
Site of malignancy	
Right lobe	56 (45.90%)
Left lobe	53 (43.44%)
Isthmus	7 (5.73%)
N/A	45 (36.88%)
Lymph Node Metastasis	
Negative	52 (42.62%)
Positive	46 (37.70%)
N/A	24 (19.68%)
Type of operation	
Total thyroidectomy and Lymph node Dissection	40 (32.78%)
Total thyroidectomy	31 (25.40%)
Right Thyroid lobectomy	8 (6.55%)
Left Thyroid lobectomy	6 (4.92%)
Partial thyroidectomy	5 (4.10%)
Completion thyroidectomy	2 (1.64%)
Completion of thyroidectomy and Lymph node Dissection	2 (1.64%)
Radio-active iodine (RAI)	
Yes	44 (36.06%)
No	22 (18.04%)
N/A	56 (45.90%)
Outcome	
No recurrence	54 (44.26%)
Recurrence	10 (8.20%)
N/A	58 (47.54%)

SD: Standard deviation, ATPO: Anti-thyroid peroxidase, NA: Not-available, TSH: Thyroid stimulating hormone, PTC: Papillary thyroid carcinoma, PTMC: Papillary thyroid microcarcinoma, MTC: Medullary thyroid carcinoma, FTC: Follicular thyroid carcinoma, FTA: Follicular thyroid adenoma, HCC: Hurthle cell carcinoma, NIFTP: Non-invasive follicular thyroid neoplasm with papillary-like nuclear features, mMTC: Medullary thyroid microcarcinoma, OC: Oncocytic carcinoma, MEC: Mucoepidermoid carcinoma, SCC: Squamous cell carcinoma, TA: Trabecular adenoma, TVC: Tall cell variant papillary thyroid carcinoma

**Table 3:** Management and outcome of collision tumors

Authors	L. Node Met.		Type of operation							RAI			Histological Findings			Site of malignancy			outcome						
	NA	+ -	TT	TT/LND	RTL	LTL	STT	CTL	CTL/LND	NA	Y	N	NA	NA	Y	N	NA	Rt. lobe		Lt. lobe	isthmus	NA	Rec	No Rec	
Awadalla et al. 2022 [13]	0	0	1	0	1	0	0	0	0	0	0	0	1	1	PTMC+FTC	0	0	0	1	PTMC+FTC	0	0	0	1	0
Abdelaal et al. 2020 [14]	0	0	6	4	0	0	1	0	0	0	0	0	6	3	PTC+FTC	3	PTMC+FTC	0	0	0	0	0	0	3	3
Abdullah et al. 2022 [1]	8	0	0	4	0	1	3	0	0	0	0	0	8	1	MTC+FTA	6	PTC+FTA	0	0	0	0	8	0	6	2
Abdullah et al. 2022 [15]	0	1	0	0	1	0	0	0	0	0	0	0	0	1	MTC+PTMC	0	1	MTC+PTMC	0	0	0	0	0	0	1
Adnan et al. 2013 [16]	0	1	3	0	3	1	0	0	0	0	0	0	4	1	PTC+MTC	2	PTC+MTC	0	0	0	0	0	0	0	4
Alavi et al. 2011 [17]	0	1	0	0	1	0	0	0	0	0	0	0	0	1	PTC+MTC	1	MTC+FTA+PTMC	0	0	0	0	0	0	0	1
Al-Mashat et al. 2003 [18]	1	0	0	0	0	1	0	0	0	0	0	0	1	1	PTC+MTC	0	1	PTC+MTC	0	0	0	0	0	0	1
Alshehri et al. 2023 [19]	0	1	0	0	1	0	0	0	0	0	0	0	1	1	PTC+MTC	0	0	0	0	0	1	0	1	0	
Polat et al. 2016 [20]	0	2	1	0	0	0	0	0	0	0	0	0	20	2	PTC+FTA	10	PTC+FTC	0	0	0	0	20	2	18	0
Baloch et al. 2001 [21]	0	1	0	0	0	0	0	0	0	0	0	0	0	1	TVC+HCC	0	1	TVC+HCC	0	0	0	0	0	0	1
Dikbas et al. 2019 [5]	0	0	1	1	0	0	0	0	0	0	0	0	1	1	PTC+MTC	1	PTC+MTC	0	0	0	0	0	0	1	0
Fallahi et al. 2023 [22]	0	3	2	0	4	0	0	0	0	1	0	0	5	1	mMTC+PTC	1	mMTC	3	1mMTC+PTC	0	2	0	0	4	0







independent origin of two tumors, proposing that their co-occurrence is a fortuitous event resulting from de novo local genesis or metastasis adjacent to an unrelated primary tumor [63,50]. Another hypothesis posits that the first tumor alters the microenvironment, thereby fostering the development of a second primary tumor. A third theory suggests that two distinct driver mutations may arise within a common stem cell, leading to the emergence of two separate tumors [64,53].

Prior studies have indicated that the predominant malignancy affecting the thyroid is PTC at 80%, succeeded by follicular thyroid carcinoma at 10%, MTC at 4%, Hürthle cell carcinoma at 3%, and anaplastic carcinoma at 2% [65]. Risk factors associated with PTC include radiation exposure, female gender, smoking, overweight or obesity, excessive dietary iodine intake, alcohol consumption, dietary nitrates, diabetes mellitus, and genetic predisposition [66].

The concurrent presence of both MTC and PTC within the same thyroid gland is a rare phenomenon, typically observed in two primary settings: a mixed tumor displaying dual differentiation or a collision tumor [17]. Rossi et al. suggested that the co-occurrence of MTC and PTC was likely coincidental. They reasoned that RET and BRAF mutations were detected in these tumors, respectively, and these mutations are independent drivers of their respective carcinomas, with a minimal chance of simultaneous occurrence [45]. Nevertheless, certain studies suggest that germline point mutations of the RET gene may play a potential role in the development of both MTC and PTC [67,68]. In the current study, a significant proportion (41.80%) of cases involving MTC and PTC as collision tumors were documented.

Adjuvant treatment, such as RAI, becomes necessary when the PTC component falls within the high or intermediate-risk category based on risk stratification. Biscolla et al. noted that the prognosis of the MTC in collision tumors remains unaffected by the presence of PTC and specific radioiodine treatments [69]. In the current study, adjuvant RAI was administered to 36.06% of the patients.

According to a review conducted by Ryan et al., until 2015, the literature documented merely 33 instances of thyroid collision tumors. Despite the notable increase in the prevalence of thyroid cancer in recent years, the simultaneous presence of multiple tumors within a single thyroid gland remains an infrequent phenomenon [1,30]. In the current study, 122 cases of thyroid collision tumors were recorded.

In a study conducted by Polat et al., 3,700 patients underwent surgery. Histopathological examination revealed benign findings in 2,686 (73%) patients and malignant findings in 1,014 (27%) patients. Among those diagnosed with differentiated thyroid carcinoma, only 20 (1.9%) had an additional neoplasm within the same thyroid gland. The average age of the patients was  $48.8 \pm 13.2$  years, with eighteen of the twenty patients being female and two being male. Lymph node metastasis was found in three patients, accounting for (15%) of the cases [7]. In the present study (71.31%) were female, while (28.69%) were male, and the mean age was  $49.87 \pm 14.48$ . However (37.70%) patients were found to have lymph node metastasis.

In another study conducted by Abdullah et al., eight cases of collision tumors were included; four patients (50%) underwent lobectomy, while the remaining four patients underwent total thyroidectomy. These cases presented various symptoms, such as neck swelling, dyspnea, and dizziness [1]. In the current study, neck swelling was the most frequently reported symptom (45.08%). Nevertheless, most of the patients (32.78%) underwent total thyroidectomy and lymph node dissection.

## 5. Conclusion

Thyroid collision tumors may primarily affect females, with the most frequent collisions being papillary and medullary thyroid carcinoma. Total thyroidectomy with lymph node dissection is the prevalent management option, and the recurrence rate can be lower than 10%.

## Declarations

**Conflicts of interest:** The author(s) have no conflicts of interest to disclose.

**Ethical approval:** Not applicable, as systematic reviews do not require ethical approval.

**Patient consent (participation and publication):** Not applicable.

**Funding:** The present study received no financial support.

**Acknowledgements:** None to be declared.

**Use of AI:** AI was not used in the drafting of the manuscript, the production of graphical elements, or the collection and analysis of data.

**Authors' contributions:** BAA and FHK were major contributors to the conception of the study, as well as to the literature search for related studies. HAN, AMS, HOB, and AMA were involved in the literature review, manuscript writing, and data analysis and interpretation. YMM, AJQ, and ASM Literature review, final approval of the manuscript, and processing of the tables. RMA, RMA, SSO, YAS, HMD and ROM were involved in the literature review, the study's design, and the manuscript's critical revision. FHK and HOA Confirmation of the authenticity of all the raw data. All authors approved the final version of the manuscript.

**Data availability statement:** Note applicable.

## References

1. Abdullah AM, Qaradakh AJ, Ahmed MM, Salih AM, Omar SS, Kakamad FH, et al. Thyroid collision tumors; A case series with literature review. *Annals of Medicine and Surgery*. 2022; 76:103444. doi:10.1016/j.amsu.2022.103444
2. Fatima ZH, Rafey M, Fazal N, Akhtar K. Synchronous cribriform morular variant of papillary and follicular carcinoma thyroid: A case series of a rare collision tumor. *Annals of Oncology Research and Therapy*. 2022; 2(2):92-6. doi:10.4103/aort.aort\_18\_22

3. Brandwein-Gensler M, Urken M, Wang B. Collision tumor of the thyroid: a case report of metastatic liposarcoma plus papillary thyroid carcinoma. *Head & Neck: Journal for the Sciences and Specialties of the Head and Neck*. 2004 ;7(7):637-41. doi: 10.1002/hed.20024
4. Ateşpare A, ÇALIŞ AB, Çelik Ö, Yener N, Vural Ç. Concurrent medullary and papillary carcinoma of thyroid. *The Turkish Journal of Ear Nose and Throat*. 2015 ;25(3):170-3. doi: 10.5606/kbbihtisas.2015.59002
5. Dikbas O, Duman AA, Guvendi GF. Medullary thyroid carcinoma and papillary thyroid carcinoma in the same patient as a collision tumour. *Case Reports in Endocrinology*. 2019 ;2019. doi:10.1155/2019/4038628
6. Sirbu AM, Sirbu CA, Eftimie L, Soare AM, Ghinescu MC, Ionita-Radu F. Multiple sclerosis, human herpesvirus 4 and thyroid collision tumor: a case report. *Experimental and Therapeutic Medicine*. 2020 ;20(4):3458-61. doi:10.3892/etm.2020.8975
7. Polat S, Arpacı D, Yazgan A, Baser H, Kilic M, Ersoy R, Kacir B. Simultaneous occurrence of different follicular neoplasms within the same thyroid gland. *Turkish Journal of Endocrinology and Metabolism*. 2016;20(2). doi:10.4274/tjem.3086
8. Jameson JL. Minimizing unnecessary surgery for thyroid nodules. *New England Journal of Medicine*. 2012 ;367(8):765-7. doi:10.1056/NEJMe1205893
9. Hawasli A, Rizzo P, Khoury H, McCaffrey JL. Can fine-needle aspiration biopsy of thyroid nodule help in determining the extent of surgery in follicular and Hurthle cell neoplasm at a community teaching institution?. *The American Surgeon*. 2002 ;68(10):907-10. doi:10.1177/000313480206801017
10. Liu F, Han F, Lu L, Chang J, Bian Z, Yao J. Thyroid collision tumour with pulmonary metastases treated effectively with nedaplatin and paclitaxel chemotherapy: A case report. *Journal of International Medical Research*. 2023 ;51(11):03000605231208602. doi:10.1177/03000605231208602
11. Ryan N, Walkden G, Lazic D, Tierney P. Collision tumors of the thyroid: A case report and review of the literature. *Head & Neck*. 2015 O ;37(10):E125-9. doi:10.1002/hed.23936
12. Muhialdeen AS, Ahmed JO, Baba HO, Abdullah IY, Hassan HA, Najar KA, Mikael TM, Mustafa MQ, Mohammed DA, Omer DA, Bapir R. Kscien's list; a new strategy to discourage predatory journals and publishers (second version). *Barw Medical Journal*. 2023. doi:10.58742/bmj.v1i1.14
13. Awadalla AS, Al Saleem MA, Al Nemer AM, Ahmed A, Al Bisher E, Al Bisher HM. Collision tumor of the thyroid gland: Follicular carcinoma and papillary microcarcinoma. *Electron J Gen Med*. 2022; 19 (6): em411. doi:10.29333/ejgm/12412
14. Abdelaal A, El Ansari W, Abusabeib A, Farghaly H, Tabe AA. Simultaneous occurrence of follicular and papillary thyroid carcinomas in same thyroid lobe: a case series of six patients from Qatar. *International journal of surgery case reports*. 2020 ;73:65-70. doi:10.1016/j.ijscr.2020.06.070
15. Abdullah AM, Ali RM, Salih KM, Mohammed KK, Kakamad FH, Salih AM. Synchronous occurrence of papillary thyroid microcarcinoma, medullary thyroid carcinoma and Hashimoto thyroiditis in a single thyroid: A case report with literature review. *International Journal of Surgery Case Reports*. 2022 ;93:106888. doi:10.1016/j.ijscr.2022.106888
16. Adnan Z, Arad E, Dana J, Shendler Y, Baron E. Simultaneous occurrence of medullary and papillary thyroid microcarcinomas: a case series and review of the literature. *Journal of Medical Case Reports*. 2013 ;7:1-5. doi:10.1186/1752-1947-7-26
17. Sadat Alavi M, Azarpira N. Medullary and papillary carcinoma of the thyroid gland occurring as a collision tumor with lymph node metastasis: A case report. *Journal of medical case reports*. 2011 ;5:1-4. doi:10.1186/1752-1947-5-590
18. Al-Mashat FM. Simultaneous medullary-papillary thyroid carcinoma. *Annals of Saudi Medicine*. 2004 ;24(3):213-5. doi:10.5144/0256-4947.2004.213
19. Alshehri S, Alsarani F, Almanea L, Alharbi A. High survival simultaneous occurrence of papillary thyroid carcinoma and ectopic thyroid with medullary thyroid carcinoma; case report and literature review. *Journal of Surgical Case Reports*. 2023 ;2023(11):rjad625. doi:10.1093/jscr/rjad625
20. Polat S, Arpacı D, Yazgan A, Baser H, Kilic M, Ersoy R, Kacir B. Simultaneous occurrence of different follicular neoplasms within the same thyroid gland. *Turkish Journal of Endocrinology and Metabolism*. 2016;20(2). doi:10.4274/tjem.3086
21. Baloch ZW, Mandel S, LiVolsi VA. Combined tall cell carcinoma and Hurthle cell carcinoma (collision tumor) of the thyroid. *Archives of pathology & laboratory medicine*. 2001 ;125(4):541-3. doi:10.5858/2001-125-0541-CTCCAH
22. Fallahi P, Patrizio A, Stoppini G, Elia G, Ragusa F, Paparo SR, Balestri E, Mazzi V, Botrini C, Varricchi G, Ulisse S. Simultaneous Occurrence of Medullary Thyroid Carcinoma and Papillary Thyroid Carcinoma: A Case Series with Literature Review. *Current Oncology*. 2023 ;30(12):10237-48. doi:10.3390/curroncol30120745
23. Fulciniti F, Vuttariello E, Calise C, Monaco M, Pezzullo L, Chiofalo MG, Di Gennaro F, Malzone MG, Campanile AC, Losito NS, Botti G. Combined papillary and mucoepidermoid carcinoma of the thyroid gland: a possible collision tumor diagnosed on fine-needle cytology. Report of a case with immunocytochemical and molecular correlations. *Endocrine pathology*. 2015 ;26:140-4. doi:10.1007/s12022-015-9364-9
24. Giacomelli L, Guerriero G, Falvo L, Altomare V, Chiesa C, Ferri S, Stio F. Simultaneous occurrence of medullary carcinoma and papillary microcarcinoma of thyroid in a patient with MEN 2A syndrome. Report of a case. *Tumori Journal*. 2007 ;93(1):109-11. doi:10.1177/030089160709300121
25. Gosavi RS, Gupta AD, Gosavi SD, Patwardhan MH. An Approach to a Rare Case of Collision Tumor of Follicular and Papillary Carcinomas of the Thyroid. doi:10.5005/jp-journals-10003-1353
26. He Q, Zhuang D, Zheng L, Zhou P, Duan S, Jing H. Collision malignant melanoma and medullary carcinoma of the thyroid. *Surgical Practice*. 2014 ;18(4):187-90. doi:10.1111/1744-1633.12090
27. He X, Soleimanpour SA, Clines GA. Adrenal metastasis as the initial diagnosis of synchronous papillary and follicular thyroid cancer. *Clinical Diabetes and Endocrinology*. 2020 ;6:1-6. doi:10.1186/s40842-020-00109-0
28. Hidangmayum A, Mathias M, HL KP, Sajitha K, Joshi D. Synchronous follicular carcinoma and non-invasive follicular thyroid neoplasm with papillary like nuclear features in thyroid-FA case report. *Biomedicine*. 2022 ;42(1):195-8. doi:10.51248/v42i1.584
29. Kawasaki K, Kai K, Tanaka N, Kido S, Ibi A, Minesaki A, Yamauchi M, Kuratomi Y, Aishima S, Nakashima M, Ito M. Collision tumor of a papillary and follicular thyroid carcinoma: a case report. *Thyroid Research*. 2023 ;16(1):24. doi:10.1186/s13044-023-00167-3
30. Kim KJ, Hong SW, Kim SM, Lee YS, Chang HS, Park CS. Synchronous occurrence of papillary, follicular, and medullary carcinoma in the same thyroid gland. doi:10.16956%2Fkjes.2014.14.3.167
31. Kobayashi K, Teramoto S, Maeta H, Ishiguro S, Mori T, Horie Y. Simultaneous occurrence of medullary carcinoma and papillary carcinoma and papillary carcinoma of the thyroid. *Journal of surgical oncology*. 1995 Aug;59(4):276-9. doi:10.1002/jso.2930590416
32. Kösem M, Kotan Ç, Algün E, Topal C. Simultaneous occurrence of papillary intrafollicular and microcarcinomas with bilateral medullary microcarcinoma of the thyroid in a patient with multiple endocrine neoplasia type 2A: report of a case. *Surgery today*. 2002 ;32:623-8. doi:10.1007/s005950200112
33. Koufopoulos N, Zacharotou A, Gouloumis AR, Papadimitriou N, Tomos P, Foukas PG, Panayiotides IG. Metastatic thyroid osteosarcoma with concomitant multifocal papillary carcinoma presenting as a collision tumor. *Cureus*. 2021 ;13(6). doi: 10.7759/cureus.15425
34. Lalanda R, Aparicio D, Boavida J, Presa DL, Batista L, Miranda L. Synchronous mixed medullary-papillary carcinoma and papillary multifocal carcinoma of the thyroid with cervical lymph node metastases. *Clinical Case Reports*. 2023 ;11(6). doi:10.1002%2Fccr3.7433
35. Lax S, Beham A, Denk H, Kronberger-Schönecker D, Langsteger W. Coexistence of papillary and medullary carcinoma of the thyroid gland-mixed or collision tumour? Clinicopathological analysis of three cases. *Virchows Archiv*. 1994 ;424:441-7. doi:10.1007/BF00190568
36. Li H, Livneh N, Dogan S, Shaha AR. The growth kinetics of collision nodal metastasis from medullary and papillary thyroid carcinomas: a case report. *European Thyroid Journal*. 2021 ;10(4):345-52. doi:10.1159/000511184
37. Ma T, Wang R, Zhou X, Liu L, Pan A, Wang H, Huang L. Case reports of collision and composite carcinomas of the thyroid: an insight into their origin and clinical significance. *BMC Endocrine Disorders*. 2023 ;23(1):173. doi:10.1186/s12902-023-01409-z
38. Mohammadzadeh M, Pourzand A, Eftekhari-Sadat A, Alikhah H, Naghavi-Behzad M. A case of concurrent several forms of thyroid cancer. *Nigerian Medical Journal*. 2013;54(5):351-3. doi:10.4103/0300-1652.122372
39. Nabili V, Natarajan S, Hirschowitz S, Bhuta S, Abemayor E. Collision tumor of thyroid: metastatic lung adenocarcinoma plus papillary thyroid carcinoma. *American journal of otolaryngology*. 2007 ;28(3):218-20. doi:10.1016/j.amjoto.2006.08.002
40. Pishdad R, Cespedes L, Boutin R, Jaloudi M, Raghuvanshi M. Coexistence of two different thyroid malignancies: a collision phenomenon. *Cureus*. 2020 ;12(4). doi:10.7759/cureus.7539
41. Plauche V, Dewenter T, Walvekar RR. Follicular and papillary carcinoma: a thyroid collision tumor. *Indian Journal of Otolaryngology and Head & Neck Surgery*. 2013 ;65:182-4. doi:10.1007/s12070-011-0450-0
42. Rachida B, Safa J, Bouaziz N, Amel EK, Mehdi F. Simultaneous medullary and papillary thyroid carcinoma as-associated with a primary renal carcinoma:

- Case report and review of literature. *J Clin Images Med Case Rep.* 2023;4(10):2636. doi: [doi:10.52768/2766-7820/2636](https://doi.org/10.52768/2766-7820/2636)
43. Rana C, Kumari N. Hurthle cell adenoma and papillary microcarcinoma in thyroid: Collision tumors. *World J Surg Oncol.* 2018 ;10(2):134-6. doi: [doi:10.5005/jp-journals-10002-1232](https://doi.org/10.5005/jp-journals-10002-1232)
  44. Rekhi B, Badhe RR, Desouza MA, Chaukar D, D'Cruz AK, Arya S, Kane SV. A unique RET EXON 11 (G691S) polymorphism in an Indian patient with a collision tumor of the thyroid. *Diagnostic Pathology.* 2007 ;2:1-8. doi: [doi:10.1186/1746-1596-2-39](https://doi.org/10.1186/1746-1596-2-39)
  45. Rossi S, Fugazzola L, De Pasquale L, Braidotti P, Cirello V, Beck-Peccoz P, Bosari S, Bastagli A. Medullary and papillary carcinoma of the thyroid gland occurring as a collision tumour: report of three cases with molecular analysis and review of the literature. *Endocrine-Related Cancer.* 2005 ;12(2):281-9. doi: [doi:10.1677/erc.1.00901](https://doi.org/10.1677/erc.1.00901)
  46. Saharan S, Solanki R, Saharan D. A rare case of collision tumor of thyroid gland: simultaneous occurrence of calcitonin negative medullary thyroid carcinoma and papillary thyroid carcinoma. *International Journal of Otorhinolaryngology and Head and Neck Surgery.* 2019 ;5(6):1726-8. doi: [doi:10.18203/issn.2454-5929.ijohns20194960](https://doi.org/10.18203/issn.2454-5929.ijohns20194960)
  47. SamieeRad F, Emami A. Synchronous occurrence of papillary thyroid carcinoma and medullary carcinoma in the setting of Hashimoto's thyroiditis and multi nodular goiter. *Iranian Journal of Pathology.* 2022;17(1):91. doi: [doi:10.306999%2FIJJP.2021.527288.2606](https://doi.org/10.306999%2FIJJP.2021.527288.2606)
  48. Stenman A, Kjellman M, Zedenius J, Juhlin CC. Synchronous lateral lymph node metastases from papillary and follicular thyroid carcinoma: case report and review of the literature. *Thyroid Research.* 2022 ;15(1):1. doi: [doi:10.1186/s13044-022-00120-w](https://doi.org/10.1186/s13044-022-00120-w)
  49. TAŞCI Hİ, Erinanc H, Emin TU, KARAGÜLLE E. Coexistence of Papillary Microcarcinoma and Hurthle Cell Adenoma: a Case of Thyroid Collision Tumor. *Clinical and Experimental Health Sciences.* 2022 ;12(4):1046-8. doi: [doi:10.33808/clinexphealthsci.938698](https://doi.org/10.33808/clinexphealthsci.938698)
  50. Tanaka T, Yoshimi N, Kanai N, Mori H, Nagai K, Fujii A, Sakata S, Tokimitsu N. Simultaneous occurrence of medullary and follicular carcinoma in the same thyroid lobe. *Human pathology.* 1989 ;20(1):83-6. doi: [doi:10.1016/0046-8177\(89\)90208-6](https://doi.org/10.1016/0046-8177(89)90208-6)
  51. Tarui T, Ishikawa N, Kadoya S, Watanabe G. Co-occurrence of papillary thyroid cancer and MALT lymphoma of the thyroid with severe airway obstruction: A case report and review of the literature. *International Journal of Surgery Case Reports.* 2014 ;5(9):594-7. doi: [doi:10.1016/j.ijscr.2014.05.016](https://doi.org/10.1016/j.ijscr.2014.05.016)
  52. Thomas VP, George R. Collision tumors of the thyroid: Review of literature and report of a case of papillary-Follicular collision tumor. *Thyroid Research and Practice.* 2018 ;15(2):60-4. doi: [doi:10.4103/trp.trp\\_6\\_18](https://doi.org/10.4103/trp.trp_6_18)
  53. Thomas A, Mittal N, Rane SU, Bal M, Patil A, Ankathi SK, Vaish R. Papillary and medullary thyroid carcinomas presenting as collision tumors: a case series of 21 cases at a tertiary care cancer center. *Head and Neck Pathology.* 2021 ;15:1137-46. doi: [doi:10.1007/s12105-021-01323-7](https://doi.org/10.1007/s12105-021-01323-7)
  54. Toyoshima MT, Domingues RB, Soares IC, Danilovic DL, Amorim LC, Cavalcante ER, Antonacio FF, Roitberg FS, Hoff AO. Thyroid collision tumor containing oncocytic carcinoma, classical and hobnail variants of papillary carcinoma and areas of poorly differentiated carcinoma. *Archives of Endocrinology and Metabolism.* 2021 ;65:495-9. doi: [doi:10.20945/2359-3997000000389](https://doi.org/10.20945/2359-3997000000389)
  55. Van Vlaenderen J, Logghe K, Schiettecatte E, Vermeersch H, Huvenne W, De Waele K, Van Beveren H, Van Dorpe J, Creyten D, De Schepper J. A synchronous papillary and follicular thyroid carcinoma presenting as a large toxic nodule in a female adolescent. *International Journal of Pediatric Endocrinology.* 2020 ;2020:1-6. doi: [doi:10.1186/s13633-020-00084-4](https://doi.org/10.1186/s13633-020-00084-4)
  56. Vlad M, Corlan A, Balas M, Golu I, Amzar D, Bistriean E, Cornianu M. Collision tumor of the thyroid—a challenge during the COVID-19 pandemic. *Archive of Clinical Cases.* 2021;8(4):64. doi: [doi:10.22551%2F2021.33.0804.10189](https://doi.org/10.22551%2F2021.33.0804.10189)
  57. Walvekar RR, Kane SV, D'Cruz AK. Collision tumor of the thyroid: follicular variant of papillary carcinoma and squamous carcinoma. *World journal of surgical oncology.* 2006 ;4:1-6. doi: [doi:10.1186/1477-7819-4-65](https://doi.org/10.1186/1477-7819-4-65)
  58. Warman M, Lipschitz N, Ikher S, Halperin D. Collision tumor of the thyroid gland: primary squamous cell and papillary thyroid carcinoma. *International Scholarly Research Notices.* 2011;2011. doi: [doi:10.5402/2011/582374](https://doi.org/10.5402/2011/582374)
  59. Whitehouse LL, Chengot P. Simultaneous papillary carcinoma and mucosa-associated lymphoid tissue (MALT) lymphoma of the thyroid; a case report. *Diagnostic Histopathology.* 2020 ;26(5):233-6. doi: [doi:10.1016/j.mpdhp.2020.02.006](https://doi.org/10.1016/j.mpdhp.2020.02.006)
  60. Younes N, Shomaf M, Al Hassan L. Simultaneous medullary and papillary thyroid carcinoma with lymph node metastasis in the same patient: case report and review of the literature. *Asian Journal of Surgery.* 2005 ;28(3):223-6. doi: [doi:10.1016/S1015-9584\(09\)60348-1](https://doi.org/10.1016/S1015-9584(09)60348-1)
  61. Ziaolhagh R, Sadrizadeh A, Shabany BP, Roudi AA. A case report of simultaneous medullary and papillary carcinoma of thyroid. *Endocrine regulations.* 2021 ;55(2):83-8. doi: [doi:10.2478/enr-2021-0010](https://doi.org/10.2478/enr-2021-0010)
  62. Sung CT, Shetty A, Menias CO, Houshyar R, Chatterjee S, Lee TK, Tung P, Helmy M, Lall C. Collision and composite tumors; radiologic and pathologic correlation. *Abdominal Radiology.* 2017 ;42:2909-26. doi: [doi:10.1007/s00261-017-1200-x](https://doi.org/10.1007/s00261-017-1200-x)
  63. Lamberg BA, Reissel P, Stenman S, Koivuniemi A, Ekblom M, Mäkinen J, Franssila K. Concurrent medullary and papillary thyroid carcinoma in the same thyroid lobe and in siblings. *Acta Medica Scandinavica.* 1981 ;209(1-6):421-4. doi: [doi:10.1111/j.0954-6820.1981.tb11620.x](https://doi.org/10.1111/j.0954-6820.1981.tb11620.x)
  64. Pastolero GC, Coire CI, Asa SL. Concurrent medullary and papillary carcinomas of thyroid with lymph node metastases: a collision phenomenon. *The American journal of surgical pathology.* 1996 ;20(2):245-50. doi: [doi:10.1089/thy.2008.0280](https://doi.org/10.1089/thy.2008.0280)
  65. Davies L, Welch HG. Increasing incidence of thyroid cancer in the United States, 1973-2002. *Jama.* 2006 ;295(18):2164-7. doi: [doi:10.1001/jama.295.18.2164](https://doi.org/10.1001/jama.295.18.2164)
  66. Sulaieva O, Selezniyov O, Shapochka D, Belemets N, Nechay O, Cheresheva Y, Tsomartova D, Ivanova M. Hashimoto's thyroiditis attenuates progression of papillary thyroid carcinoma: deciphering immunological links. *Heliyon.* 2020 ;6(1). doi: [doi:10.1016/j.heliyon.2019.e03077](https://doi.org/10.1016/j.heliyon.2019.e03077)
  67. Brauckhoff M, Gimm O, Hinze R, Ukkat J, Brauckhoff K, Dralle H. Papillary thyroid carcinoma in patients with RET proto-oncogene germline mutation. *Thyroid.* 2002 ;12(7):557-61. doi: [doi:10.1089/105072502320288393](https://doi.org/10.1089/105072502320288393)
  68. Papi G, Corrado S, Pomponi MG, Carapezzi C, Cesinaro A, LiVolsi VA. Concurrent lymph node metastases of medullary and papillary thyroid carcinoma in a case with RET oncogene germline mutation. *Endocrine Pathology.* 2003 ;14:269-76. doi: [doi:10.1007/s12022-003-0020-4](https://doi.org/10.1007/s12022-003-0020-4)
  69. Pelizzo MR, Bernante P, Toniato A, Fassina A. Frequency of thyroid carcinoma in a recent series of 539 consecutive thyroidectomies for multinodular goiter. *Tumori Journal.* 1997 ;83(3):653-5. doi: [doi:10.1177/030089169708300305](https://doi.org/10.1177/030089169708300305)